



The Tall Wind project - exploring the wind profile and boundary-layer height in the atmosphere's first kilometer over flat terrain

Gryning, Sven-Erik; Batchvarova, Ekaterina; Pena Diaz, Alfredo; Mikkelsen, Torben; Brümmer, B.; Emeis, S.; Gulstad, L.; Lee, N.

Published in:
Proceedings

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Gryning, S-E., Batchvarova, E., Pena Diaz, A., Mikkelsen, T., Brümmer, B., Emeis, S., Gulstad, L., & Lee, N. (2010). The Tall Wind project - exploring the wind profile and boundary-layer height in the atmosphere's first kilometer over flat terrain. In *Proceedings*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The Tall Wind project – exploring the wind profile and boundary-layer height in the atmosphere's first kilometer over flat terrain.

S.E. Gryning¹, E. Batchvarova¹, A. Pena¹, T. Mikkelsen¹,
B. Brümmer², S. Emeis³, L. Gulstad⁴, and N. Lee⁵


¹Risø DTU, ²Hamburg University, ³Karlsruhe Institute of
Technology, ⁴Vestas A/S, ⁵DONG Energy

Measuring sites




The study of the wind profile over flat homogeneous terrain in The Tall Wind project is based on a combination of measurements, data analysis and modelling. The idea is to supplement existing long-term measurements of turbulence and wind speed at tall well instrumented meteorological masts (at Høvsøre to reflect low roughness over rural area; and at Hamburg to reflect high roughness such as urban conditions) and at Horns Rev 2 (off-shore, not instrumented with turbulence instruments). Additional measurements will address the wind profile up to 500 to 1000 meters by use of a newly developed commercial pulsed wind Lidar (WSL70). The height of the planetary boundary layer will be determined from aerosol backscatter with a new meteorological Lidar (ASL 300).



Aerosol lidar



Wind lidar



Short term averaged wind profile (i. e. 0.5 hour)




$$U = \frac{u_{*o}}{\kappa} \left[\ln\left(\frac{z}{z_o}\right) + \frac{z}{l_{\text{MBL}}} - \frac{z}{z_i} \frac{z}{2l_{\text{MBL}}} \right]$$

Climatological wind profile (1 month to several years)


$$\left\langle \frac{kU}{u_{*0}} \right\rangle (z) = \int \ln(z/z_0) P(z_0) dz_0 - \int \psi(z/L) P(1/L) d(1/L)$$

$$\left\langle \frac{kU}{u_{*0}} \right\rangle (z) = \ln(z/z_{0m}) - [\langle \psi \rangle (z) - \langle \psi_0 \rangle],$$

Where $\langle \psi \rangle (z)$ is the long term stability correction based on a given stability distribution. To determine the climatological wind profile we need a representation of $P(1/L)$ in measurable quantities – is worked out in Kelly and Gryning (2010).



$$\langle \psi(z/L) \rangle \neq \psi(z/\langle L \rangle)$$




$$\left\langle \frac{kU}{u_{*0}} \right\rangle (z) = \ln(z/z_{0m}) - \left[1 - \frac{z}{h_{\text{eff}}} \right] \langle \psi \rangle (z) - \frac{z}{h_{\text{eff}}} \frac{1}{z} \int_{z_0}^z \langle \psi \rangle (z) dz$$

$$+ \frac{h_{\text{eff}}}{2L_{\text{mid}}^{\text{eff}}} \left[1 - \left(1 - \frac{z}{h_{\text{eff}}} \right)^2 \right],$$

For applied use we work on the suggestion:

$$\left\langle \frac{kU}{u_{*0}} \right\rangle (z) = \ln\left(\frac{z}{z_0}\right) - \left[1 - \frac{z}{h_{\text{eff}}} \right] \langle \psi \rangle (z) - \frac{z}{h_{\text{eff}}} \langle \psi \rangle (z/2) + \left[1 - \left(1 - \frac{z}{h_{\text{eff}}} \right)^2 \right]$$

Monthly mean wind profiles at Høvsøre



Climatological wind profile

Red: unstable only

Black: all measurements

Blue: stable only

Andreas Blatt,
master thesis Risø DTU, 2010

The ultimate goal of Tall Wind Project is to come up with:

- recommendations for parameterizations for applied use of the wind profile above the surface layer
- recommendations for the measurements and instruments requirements that can provide the necessary input for yield assessments and constitute a design load basis for wind turbines.

It is realized that this is a very ambitious goal but the data set we hope to achieve will be internationally unique and an excellent basis for the research leading to better tools for the prediction of the energy yield and design basis for large wind turbines.

Thanks for your attention

ACKNOWLEDGEMENT

The study is supported by:
the Danish Council for Strategic Research,
Sagsnr. 2104-08-0025 and
the EU FP7 Marie Curie Fellowship VSABLA
(PIEF-GA-2009-237471).

